

A theory of tensor products for module categories for a vertex operator algebra, II

Yi-Zhi Huang and James Lepowsky

Contents

7	Isomorphisms between certain spaces of intertwining operators	2
8	Commutator formula for vertex operators on $(W_1 \otimes W_2)^*$	9
9	Some useful formulas	14
10	Jacobi identity for vertex operators on a certain subspace of $(W_1 \otimes W_2)^*$	16
11	Stability of certain subspaces of $(W_1 \otimes W_2)^*$ under the action of vertex operators	27

Abstract

This is the second part in a series of papers presenting a theory of tensor products for module categories for a vertex operator algebra. In Part I (hep-th/9309076), the notions of $P(z)$ - and $Q(z)$ -tensor product of two modules for a vertex operator algebra were introduced and under a certain hypothesis, two constructions of a $Q(z)$ -tensor product were given, using certain results stated without proof. In Part II, the proofs of those results are supplied.

The present paper (Part II) is a continuation of [HL] (Part I), to which the reader is referred for the necessary background, including references. In Part I, the notions of $P(z)$ - and $Q(z)$ -tensor product of two modules for a vertex operator algebra were introduced and under a certain hypothesis, two constructions of a $Q(z)$ -tensor product were given based on certain results (Proposition 4.9, Proposition 5.2, Theorem 6.1 and Proposition 6.2) stated without proof. In Part II, we supply the proofs of these results.

We fix a vertex operator algebra V and V -modules (W_1, Y_1) , (W_2, Y_2) and (W_3, Y_3) . We add $y, y_1, y_2, \dots$ to our list of commuting formal variables in Part II. The numberings of sections, formulas, etc., continue those of Part I.

Part II is organized as follows: Section 7 proves that the spaces of intertwining operators of certain types are isomorphic (Proposition 4.9). Section 8 is devoted to the proof of the commutator formula for the vertex operators defined in Section 5 (Proposition 5.2), and a related remark pointing out that the commutator formula and Proposition 5.1 in fact come from analogous properties of an action on the vector space $W_1 \otimes W_2$. In Section 9, we derive some formulas involving vertex operators and the Virasoro operators on the dual space of the tensor product vector space of two modules. In Section 10, we prove Theorem 6.1, asserting the Jacobi identity for the vertex operators acting on a certain subspace of this dual space, and in Section 11, we establish Proposition 6.2, stating the invariance of certain subspaces under these operators.

7 Isomorphisms between certain spaces of intertwining operators

In this section we prove various results, including Proposition 4.9, establishing isomorphisms between certain spaces of intertwining operators. These results are all simple generalizations of the corresponding results in [FHL], where only the monodromy-free case was considered.

Let $\mathcal{Y}$ be an intertwining operator of type $\begin{pmatrix} W_3 \\ W_1 W_2 \end{pmatrix}$. For any complex number ζ and any $w_{(1)} \in W_1$, $\mathcal{Y}(w_{(1)}, y) \Big|_{y^n = e^{n\zeta} x^n, n \in \mathbb{C}}$ is a well-defined element of $\text{Hom}(W_2, W_3)\{x\}$. We denote this element by $\mathcal{Y}(w_{(1)}, e^\zeta x)$. Note that this element depends on ζ , not on just e^ζ . Given any $r \in \mathbb{Z}$, we define

$\Omega_r(\mathcal{Y}) : W_2 \otimes W_1 \rightarrow W_3\{x\}$ by the formula

$$\Omega_r(\mathcal{Y})(w_{(2)}, x)w_{(1)} = e^{xL(-1)}\mathcal{Y}(w_{(1)}, e^{(2r+1)\pi i}x)w_{(2)}. \quad (7.1)$$

for $w_{(1)} \in W_1$ and $w_{(2)} \in W_2$.

Proposition 7.1 *The operator $\Omega_r(\mathcal{Y})$ is an intertwining operator of type $\begin{pmatrix} W_3 \\ W_2 \ W_1 \end{pmatrix}$. Moreover,*

$$\Omega_{-r-1}(\Omega_r(\mathcal{Y})) = \Omega_r(\Omega_{-r-1}(\mathcal{Y})) = \mathcal{Y}. \quad (7.2)$$

In particular, the correspondence $\mathcal{Y} \mapsto \Omega_r(\mathcal{Y})$ defines a linear isomorphism from $\mathcal{V}_{W_1 W_2}^{W_3}$ to $\mathcal{V}_{W_2 W_1}^{W_3}$, and we have

$$N_{W_1 W_2}^{W_3} = N_{W_2 W_1}^{W_3}. \quad (7.3)$$

Proof The proof of this result is a straightforward generalization of the proof of Proposition 5.4.7 in [FHL]. The lower truncation condition (2.44) is clear. From the Jacobi identity

$$\begin{aligned} & x_0^{-1}\delta\left(\frac{x_1 - y}{x_0}\right)Y_3(v, x_1)\mathcal{Y}(w_{(1)}, y)w_{(2)} \\ & \quad - x_0^{-1}\delta\left(\frac{y - x_1}{-x_0}\right)\mathcal{Y}(w_{(1)}, y)Y_2(v, x_1)w_{(2)} \\ & = y^{-1}\delta\left(\frac{x_1 - x_0}{y}\right)\mathcal{Y}(Y_1(v, x_0)w_{(1)}, y)w_{(2)} \end{aligned} \quad (7.4)$$

for $\mathcal{Y}$, we obtain

$$\begin{aligned} & x_0^{-1}\delta\left(\frac{x_1 - y}{x_0}\right)e^{-yL(-1)}Y_3(v, x_1)\mathcal{Y}(w_{(1)}, y)w_{(2)}\Big|_{y^n = e^{n(2r+1)\pi i}x_2^n, n \in \mathbb{C}} \\ & \quad - x_0^{-1}\delta\left(\frac{y - x_1}{-x_0}\right)e^{-yL(-1)}\mathcal{Y}(w_{(1)}, y)Y_2(v, x_1)w_{(2)}\Big|_{y^n = e^{n(2r+1)\pi i}x_2^n, n \in \mathbb{C}} \\ & = y^{-1}\delta\left(\frac{x_1 - x_0}{y}\right)e^{-yL(-1)}\mathcal{Y}(Y_1(v, x_0)w_{(1)}, y)w_{(2)}\Big|_{y^n = e^{n(2r+1)\pi i}x_2^n, n \in \mathbb{C}}. \end{aligned} \quad (7.5)$$

The first term of the left-hand side of (7.5) is equal to

$$\begin{aligned} & x_0^{-1}\delta\left(\frac{x_1 - y}{x_0}\right)Y_3(v, x_1 - y)e^{-yL(-1)}\mathcal{Y}(w_{(1)}, y)w_{(2)}\Big|_{y^n = e^{n(2r+1)\pi i}x_2^n, n \in \mathbb{C}} \\ & = x_0^{-1}\delta\left(\frac{x_1 + x_2}{x_0}\right)Y_3(v, x_0)\Omega_r(\mathcal{Y})(w_{(2)}, x_2)w_{(1)}, \end{aligned}$$

the second term is equal to

$$-x_0^{-1} \delta \left(\frac{-x_2 - x_1}{-x_0} \right) \Omega_r(\mathcal{Y})(Y_2(v, x_1)w_{(2)}, x_2)w_{(1)}$$

and the right-hand side of (7.5) is equal to

$$-x_2^{-1} \delta \left(\frac{x_1 - x_0}{-x_2} \right) \Omega_r(\mathcal{Y})(w_{(2)}, x_2)Y_1(v, x_0)w_{(1)}.$$

Substituting into (7.5) we obtain

$$\begin{aligned} & x_0^{-1} \delta \left(\frac{x_1 + x_2}{x_0} \right) Y_3(v, x_0) \Omega_r(\mathcal{Y})(w_{(2)}, x_2)w_{(1)} \\ & \quad - x_0^{-1} \delta \left(\frac{x_2 + x_1}{x_0} \right) \Omega_r(\mathcal{Y})(Y_2(v, x_1)w_{(2)}, x_2)w_{(1)} \\ & = -x_2^{-1} \delta \left(\frac{x_1 - x_0}{-x_2} \right) \Omega_r(\mathcal{Y})(w_{(2)}, x_2)Y_1(v, x_0)w_{(1)}, \end{aligned} \quad (7.6)$$

which by (2.6) is equivalent to

$$\begin{aligned} & x_1^{-1} \delta \left(\frac{x_0 - x_2}{x_1} \right) Y_3(v, x_0) \Omega_r(\mathcal{Y})(w_{(2)}, x_2)w_{(1)} \\ & \quad - x_1^{-1} \delta \left(\frac{x_2 - x_0}{-x_1} \right) \Omega_r(\mathcal{Y})(w_{(2)}, x_2)Y_1(v, x_0)w_{(1)} \\ & = x_2^{-1} \delta \left(\frac{x_0 - x_1}{x_2} \right) \Omega_r(\mathcal{Y})(Y_2(v, x_1)w_{(2)}, x_2)w_{(1)}. \end{aligned} \quad (7.7)$$

This in turn is the Jacobi identity for $\Omega_r(\mathcal{Y})$. The $L(-1)$ -derivative property of $\Omega_r(\mathcal{Y})$ is easily proved as in the proof of Proposition 5.4.7 of [FHL]. The identity (7.2) is obvious from the definitions of $\Omega_r(\mathcal{Y})$ and $\Omega_{-r-1}(\mathcal{Y})$. $\square$

Remark 7.2 As Propositions 4.7 and 7.1 suggest, for each triple $s_1, s_2, s_3 \in \mathbb{Z}$ the intertwining operator $\mathcal{Y}$ gives rise to an intertwining operator $\mathcal{Y}_{[s_1, s_2, s_3]}$ of the same type, defined by

$$\mathcal{Y}_{[s_1, s_2, s_3]}(w_{(1)}, x) = e^{2s_1\pi i L(0)} \mathcal{Y}(e^{2s_2\pi i L(0)} w_{(1)}, x) e^{2s_3\pi i L(0)} \quad (7.8)$$

for $w_{(1)} \in W_1$; it is easy to see that $\mathcal{Y}_{[s_1, s_2, s_3]}$ is again an intertwining operator. We have $\mathcal{Y}_{[0, 0, 0]} = \mathcal{Y}$ and for $r_1, r_2, r_3, s_1, s_2, s_3 \in \mathbb{Z}$,

$$(\mathcal{Y}_{[r_1, r_2, r_3]})_{[s_1, s_2, s_3]} = \mathcal{Y}_{[r_1 + s_1, r_2 + s_2, r_3 + s_3]}. \quad (7.9)$$

For any $a \in \mathbb{C}$, we have the formula

$$e^{aL(0)} \mathcal{Y}(w_{(1)}, x) e^{-aL(0)} = \mathcal{Y}(e^{aL(0)} w_{(1)}, e^a x) \quad (7.10)$$

whose proof is similar to that of formula (5.4.22) of [FHL]. From this formula, we see that (7.2) generalizes to:

$$\begin{aligned} \Omega_s(\Omega_r(\mathcal{Y})) &= \mathcal{Y}_{[r+s+1, -(r+s+1), -(r+s+1)]} \\ &= \mathcal{Y}(\cdot, e^{2(r+s+1)\pi i} \cdot). \end{aligned} \quad (7.11)$$

Given an intertwining operator $\mathcal{Y}$ of type $\begin{pmatrix} W_3 \\ W_1 W_2 \end{pmatrix}$ and an integer $r \in \mathbb{Z}$, we define the r -contragredient operator of $\mathcal{Y}$ to be the linear map

$$\begin{aligned} W_1 \otimes W'_3 &\rightarrow W'_2\{z\} \\ w_{(1)} \otimes w'_{(3)} &\mapsto A_r(\mathcal{Y})(w_{(1)}, x) w'_{(3)} \end{aligned} \quad (7.12)$$

given by

$$\begin{aligned} \langle A_r(\mathcal{Y})(w_{(1)}, x) w'_{(3)}, w_{(2)} \rangle_{W_2} &= \\ = \langle w'_{(3)}, \mathcal{Y}(e^{xL(1)} e^{(2r+1)\pi i L(0)} x^{-2L(0)} w_{(1)}, x^{-1}) w_{(2)} \rangle_{W_3}, \end{aligned} \quad (7.13)$$

where $w_1 \in W_1$, $w_2 \in W_2$, $w'_{(3)} \in W'_3$ and $\langle \cdot, \cdot \rangle_{W_2}$ and $\langle \cdot, \cdot \rangle_{W_3}$ are the pairings for W'_2, W_2 and for W'_3, W_3 , respectively. Note that for the case $W_1 = V$ and $W_2 = W_3 = W$, the operator $A_r(\mathcal{Y})$ agrees with the contragredient vertex operator $\mathcal{Y}'$ (recall (2.48), (2.49)) for any $r \in \mathbb{Z}$.

Proposition 7.3 *The r -contragredient operator $A_r(\mathcal{Y})$ of an intertwining operator $\mathcal{Y}$ of type $\begin{pmatrix} W_3 \\ W_1 W_2 \end{pmatrix}$ is an intertwining operator of type $\begin{pmatrix} W'_2 \\ W_1 W'_3 \end{pmatrix}$. Moreover,*

$$A_{-r-1}(A_r(\mathcal{Y})) = A_r(A_{-r-1}(\mathcal{Y})) = \mathcal{Y}. \quad (7.14)$$

In particular, the correspondence $\mathcal{Y} \mapsto A_r(\mathcal{Y})$ defines a linear isomorphism from $\mathcal{V}_{W_1 W_2}^{W_3}$ to $\mathcal{V}_{W_1 W'_3}^{W'_2}$, and we have

$$N_{W_1 W_2}^{W_3} = N_{W_1 W'_3}^{W'_2}. \quad (7.15)$$

Proof The proof of this result is a straightforward generalization of the proof of Theorem 5.5.1 and Proposition 5.5.2 in [FHL]. The lower truncation condition (2.44) being easy to verify, we want to prove the identity

$$\begin{aligned}
& \langle x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y_2'(v, x_1) A_r(\mathcal{Y})(w_{(1)}, x_2) w'_{(3)}, w_{(2)} \rangle_{W_2} \\
& \quad - \langle x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) A_r(\mathcal{Y})(w_{(1)}, x_2) Y_3'(v, x_1) w'_{(3)}, w_{(2)} \rangle_{W_2} \\
& = \langle x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) A_r(\mathcal{Y})(Y_1(v, x_0) w_{(1)}, x_2) w'_{(3)}, w_{(2)} \rangle_{W_2}. \quad (7.16)
\end{aligned}$$

But by the definitions (2.49) and (7.13) we have

$$\begin{aligned}
& \langle Y_2'(v, x_1) A_r(\mathcal{Y})(w_{(1)}, x_2) w'_{(3)}, w_{(2)} \rangle_{W_2} = \\
& = \langle w'_{(3)}, \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) \cdot \\
& \quad \cdot Y_2(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, x_1^{-1}) w_{(2)} \rangle_{W_3}, \quad (7.17)
\end{aligned}$$

$$\begin{aligned}
& \langle A_r(\mathcal{Y})(w_{(1)}, x_2) Y_3'(v, x_1) w'_{(3)}, w_{(2)} \rangle_{W_2} = \\
& = \langle w'_{(3)}, Y_3(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, x_1^{-1}) \cdot \\
& \quad \cdot \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) w_{(2)} \rangle_{W_3}, \quad (7.18)
\end{aligned}$$

$$\begin{aligned}
& \langle A_r(\mathcal{Y})(Y_1(v, x_0) w_{(1)}, x_2) w'_{(3)}, w_{(2)} \rangle_{W_2} = \\
& = \langle w'_{(3)}, \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} Y_1(v, x_0) w_{(1)}, x_2^{-1}) w_{(2)} \rangle_{W_3}, \quad (7.19)
\end{aligned}$$

and from the Jacobi identity for $\mathcal{Y}$ we have

$$\begin{aligned}
& \langle w'_{(3)}, \left(\frac{-x_0}{x_1 x_2} \right)^{-1} \delta \left(\frac{x_1^{-1} - x_2^{-1}}{-x_0/x_1 x_2} \right) Y_3(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, x_1^{-1}) \cdot \\
& \quad \cdot \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) w_{(2)} \rangle_{W_3} \\
& - \langle w'_{(3)}, \left(\frac{-x_0}{x_1 x_2} \right)^{-1} \delta \left(\frac{x_2^{-1} - x_1^{-1}}{x_0/x_1 x_2} \right) \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) \cdot \\
& \quad \cdot Y_2(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, x_1^{-1}) w_{(2)} \rangle_{W_3} \\
& = \langle w'_{(3)}, (x_2^{-1})^{-1} \delta \left(\frac{x_1^{-1} + x_0/x_1 x_2}{x_2^{-1}} \right) \mathcal{Y}(Y_1(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, -x_0/x_1 x_2) \cdot \\
& \quad \cdot e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) w_{(2)} \rangle_{W_3}, \quad (7.20)
\end{aligned}$$

or equivalently,

$$\begin{aligned}
& -\langle w'_{(3)}, x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) Y_3(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, x_1^{-1}) \cdot \\
& \quad \cdot \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) w_{(2)} \rangle_{W_3} \\
& + \langle w'_{(3)}, x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) \cdot \\
& \quad \cdot Y_2(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, x_1^{-1}) w_{(2)} \rangle_{W_3} \\
& = \langle w'_{(3)}, x_1^{-1} \delta \left(\frac{x_2 + x_0}{x_1} \right) \mathcal{Y}(Y_1(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, -x_0/x_1 x_2) \cdot \\
& \quad \cdot e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}) w_{(2)} \rangle_{W_3}. \tag{7.21}
\end{aligned}$$

Substituting (7.17), (7.18) and (7.19) into (7.16) and then comparing with (7.21), we see that the proof of (7.16) is reduced to the proof of

$$\begin{aligned}
& x_1^{-1} \delta \left(\frac{x_2 + x_0}{x_1} \right) \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} Y_1(v, x_0) w_{(1)}, x_2^{-1}) \\
& = x_1^{-1} \delta \left(\frac{x_2 + x_0}{x_1} \right) \mathcal{Y}(Y_1(e^{x_1 L(1)} (-x_1^{-2})^{L(0)} v, -x_0/x_1 x_2) \cdot \\
& \quad \cdot e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}), \tag{7.22}
\end{aligned}$$

or of

$$\begin{aligned}
& \mathcal{Y}(e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} Y_1(v, x_0) w_{(1)}, x_2^{-1}) = \\
& = \mathcal{Y}(Y_1(e^{(x_2+x_0)L(1)} (-(x_2+x_0)^{-2})^{L(0)} v, -x_0/(x_2+x_0)x_2) \cdot \\
& \quad \cdot e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} w_{(1)}, x_2^{-1}). \tag{7.23}
\end{aligned}$$

We see that we need only prove

$$\begin{aligned}
& e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} Y_1(v, x_0) = \\
& = Y_1(e^{(x_2+x_0)L(1)} (-(x_2+x_0)^{-2})^{L(0)} v, -x_0/(x_2+x_0)x_2) \cdot \\
& \quad \cdot e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} \tag{7.24}
\end{aligned}$$

or equivalently, the conjugation formula

$$\begin{aligned}
& e^{x_2 L(1)} e^{(2r+1)\pi i L(0)} x_2^{-2L(0)} Y_1(v, x_0) x_2^{2L(0)} e^{-(2r+1)\pi i L(0)} e^{-x_2 L(1)} \\
& = Y_1(e^{(x_2+x_0)L(1)} (-(x_2+x_0)^{-2})^{L(0)} v, -x_0/(x_2+x_0)x_2) \tag{7.25}
\end{aligned}$$

for $v \in V$, acting on the module W_1 . But formula (7.25) follows from Lemma 5.2.3 of [FHL] and the formula

$$e^{(2r+1)\pi i L(0)} Y_1(v, x) e^{-(2r+1)\pi i L(0)} = Y_1((-1)^{L(0)} v, -x), \quad (7.26)$$

which is a special case of (7.10). This establishes the Jacobi identity, and the $L(-1)$ -derivative property follows from the same argument used in the proof of Theorem 5.5.1 of [FHL]. Finally the relation (7.14) is obtained by combining the argument in the proof of Proposition 5.3.1 of [FHL] and the formula

$$e^{(2r+1)\pi i L(0)} x^{2L(0)} e^{xL(1)} x^{-2L(0)} e^{-(2r+1)\pi i L(0)} = e^{-x^{-1}L(1)} \quad (7.27)$$

on W_1 , whose proof is similar to that of formula (5.3.1) of [FHL]. $\square$

Remark 7.4 This argument shows that for any $r, s \in \mathbb{Z}$, formula (7.14) generalizes to:

$$A_s(A_r(\mathcal{Y})) = \mathcal{Y}_{[0, r+s+1, 0]}. \quad (7.28)$$

From Proposition 7.1, for any integer r_1 we have an isomorphism Ω_{r_1} from $\mathcal{V}_{W_1 W_2}^{W_3}$ to $\mathcal{V}_{W_2 W_1}^{W_3}$, and from Proposition 7.3, for any integer r_2 we have an isomorphism A_{r_2} from $\mathcal{V}_{W_2 W_1}^{W_3}$ to $\mathcal{V}_{W_2 W'_3}^{W'_1}$. By Proposition 7.1 again, for any integer r_3 there is an isomorphism, which we again denote Ω_{r_3} , from $\mathcal{V}_{W_2 W'_3}^{W'_1}$ to $\mathcal{V}_{W'_3 W_2}^{W'_1}$. Thus for any triple (r_1, r_2, r_3) of integers, we have an isomorphism $\Omega_{r_3} \circ A_{r_2} \circ \Omega_{r_1}$ from $\mathcal{V}_{W_1 W_2}^{W_3}$ to $\mathcal{V}_{W'_3 W_2}^{W'_1}$. Let $\mathcal{Y}$ be an intertwining operator in $\mathcal{V}_{W_1 W_2}^{W_3}$ and $w_{(1)}$, $w_{(2)}$, $w'_{(3)}$ elements of W_1 , W_2 , W'_3 , respectively. From the definitions of Ω_{r_1} , A_{r_2} and Ω_{r_3} , we have

$$\begin{aligned} & \langle (\Omega_{r_3} \circ A_{r_2} \circ \Omega_{r_1})(\mathcal{Y})(w'_{(3)}, x) w_{(2)}, w_{(1)} \rangle_{W_1} = \\ &= \langle e^{xL(-1)} A_{r_2}(\Omega_{r_1}(\mathcal{Y}))(w_{(2)}, e^{(2r_3+1)\pi i} x) w'_{(3)}, w_{(1)} \rangle_{W_1} \\ &= \langle A_{r_2}(\Omega_{r_1}(\mathcal{Y}))(w_{(2)}, e^{(2r_3+1)\pi i} x) w'_{(3)}, e^{xL(1)} w_{(1)} \rangle_{W_1} \\ &= \langle w'_{(3)}, \Omega_{r_1}(\mathcal{Y})(e^{-xL(1)} e^{(2r_2+1)\pi i L(0)} e^{-2(2r_3+1)\pi i L(0)} x^{-2L(0)} w_{(2)}, \\ &\quad e^{-(2r_3+1)\pi i} x^{-1}) e^{xL(1)} w_{(1)} \rangle_{W_3} \\ &= \langle w'_{(3)}, e^{-x^{-1}L(-1)} \mathcal{Y}(e^{xL(1)} w_{(1)}, e^{(2r_1+1)\pi i} e^{-(2r_3+1)\pi i} x^{-1}) \cdot \\ &\quad \cdot e^{-xL(1)} e^{(2r_2+1)\pi i L(0)} e^{-2(2r_3+1)\pi i L(0)} x^{-2L(0)} w_{(2)} \rangle_{W_3} \end{aligned}$$

$$= \langle e^{-x^{-1}L(1)}w'_{(3)}, \mathcal{Y}(e^{xL(1)}w_{(1)}, e^{2(r_1-r_3)\pi i}x^{-1}) \cdot e^{-xL(1)}e^{(2(r_2-2r_3-1)+1)\pi iL(0)}x^{-2L(0)}w_{(2)} \rangle_{W_3}. \quad (7.29)$$

From (7.29) we see that $\Omega_{r_3} \circ A_{r_2} \circ \Omega_{r_1}$ depends only on $r_2 - 2r_3 - 1$ and $r_1 - r_3$, and the operators $\Omega_{r_3} \circ A_{r_2} \circ \Omega_{r_1}$ with different $r_1 - r_3$ but the same $r_2 - 2r_3 - 1$ differ from each other only by automorphisms of $\mathcal{V}_{W_1W_2}^{W_3}$. Thus for our purpose, we need only consider those isomorphisms such that $r_1 - r_3 = 0$. Given any integer r , we choose two integers r_2 and r_3 such that $r = r_2 - 2r_3 - 1$ and we define

$$B_r = \Omega_{r_3} \circ A_{r_2} \circ \Omega_{r_3}. \quad (7.30)$$

From (7.29) we see that B_r is independent of the choices of r_2 and r_3 and that (4.31) holds. This proves Proposition 4.9.

Set

$$N_{W_1W_2W_3} = N_{W_1W_2}^{W'_3}. \quad (7.31)$$

Then from Proposition 7.1 we have

$$N_{W_1W_2W_3} = N_{W_2W_1W_3} \quad (7.32)$$

and from Proposition 7.3 we have

$$N_{W_1W_2W_3} = N_{W_1W_3W_2}, \quad (7.33)$$

so that we obtain (cf. formula (5.5.8) of [FHL]):

Proposition 7.5 *For any permutation σ of $\{1, 2, 3\}$,*

$$N_{W_1W_2W_3} = N_{W_{\sigma(1)}W_{\sigma(2)}W_{\sigma(3)}}. \quad \square \quad (7.34)$$

8 Commutator formula for vertex operators on $(W_1 \otimes W_2)^*$

We prove Proposition 5.2 in this section. The reader should note the well-definedness of each expression and the justifiability of each use of a δ -function

property in the argument which follows. Let $\lambda \in (W_1 \otimes W_2)^*$, $v_1, v_2 \in V$, $w_{(1)} \in W_1$ and $w_{(2)} \in W_2$. By (5.4),

$$\begin{aligned}
& (Y'_{Q(z)}(v_1, x_1)Y'_{Q(z)}(v_2, x_2)\lambda)(w_{(1)} \otimes w_{(2)}) = \\
& = \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) (Y'_{Q(z)}(v_2, x_2)\lambda)(Y_1^*(v_1, y_1)w_{(1)} \otimes w_{(2)}) \\
& \quad - \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) (Y'_{Q(z)}(v_2, x_2)\lambda)(w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& = \text{Res}_{y_1} \text{Res}_{y_2} x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_2, y_2)Y_1^*(v_1, y_1)w_{(1)} \otimes w_{(2)}) \\
& \quad - \text{Res}_{y_1} \text{Res}_{y_2} x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_1, y_1)w_{(1)} \otimes Y_2(v_2, y_2)w_{(2)}) \\
& \quad - \text{Res}_{y_1} \text{Res}_{y_2} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_2, y_2)w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& \quad + \text{Res}_{y_1} \text{Res}_{y_2} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, y_2)Y_2(v_1, y_1)w_{(2)}). \tag{8.1}
\end{aligned}$$

Transposing the subscripts 1 and 2 of the symbols v , x and y , we have

$$\begin{aligned}
& (Y'_{Q(z)}(v_2, x_2)Y'_{Q(z)}(v_1, x_1)\lambda)(w_{(1)} \otimes w_{(2)}) = \\
& = \text{Res}_{y_2} \text{Res}_{y_1} x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_1, y_1)Y_1^*(v_2, y_2)w_{(1)} \otimes w_{(2)}) \\
& \quad - \text{Res}_{y_2} \text{Res}_{y_1} x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_2, y_2)w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& \quad - \text{Res}_{y_2} \text{Res}_{y_1} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_1, y_1)w_{(1)} \otimes Y_2(v_2, y_2)w_{(2)}) \\
& \quad + \text{Res}_{y_2} \text{Res}_{y_1} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_1, y_1)Y_2(v_2, y_2)w_{(2)}). \tag{8.2}
\end{aligned}$$

The equalities (8.1) and (8.2) give

$$\begin{aligned}
& ([Y'_{Q(z)}(v_1, x_1), Y'_{Q(z)}(v_2, x_2)]\lambda)(w_{(1)} \otimes w_{(2)}) = \\
& = \text{Res}_{y_2} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) \cdot \\
& \quad \cdot \lambda([Y_1^*(v_2, y_2), Y_1^*(v_1, y_1)]w_{(1)} \otimes w_{(2)}) \\
& \quad - \text{Res}_{y_2} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes [Y_2(v_1, y_1), Y_2(v_2, y_2)]w_{(2)}) \\
& = \text{Res}_{y_2} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) \cdot \\
& \quad \cdot \lambda \left(\text{Res}_{x_0} y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) Y_1^*(Y(v_1, x_0)v_2, y_2)w_{(1)} \otimes w_{(2)} \right) \\
& \quad - \text{Res}_{y_2} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot \lambda \left(w_{(1)} \otimes \text{Res}_{x_0} y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) Y_2(Y(v_1, x_0)v_2, y_2)w_{(2)} \right) \\
& = \text{Res}_{x_0} \text{Res}_{y_2} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(Y(v_1, x_0)v_2, y_2)w_{(1)} \otimes w_{(2)}) \\
& \quad - \text{Res}_{x_0} \text{Res}_{y_2} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(Y(v_1, x_0)v_2, y_2)w_{(2)}). \tag{8.3}
\end{aligned}$$

Since

$$\begin{aligned}
& x_1^{-1} \delta \left(\frac{y_1 - z}{x_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) = \\
& = y_1^{-1} \delta \left(\frac{x_1 + z}{y_1} \right) y_2^{-1} \delta \left(\frac{x_2 + z}{y_2} \right) (x_2 + z)^{-1} \delta \left(\frac{(x_1 + z) - x_0}{x_2 + z} \right) \\
& = y_1^{-1} \delta \left(\frac{x_1 + z}{y_1} \right) y_2^{-1} \delta \left(\frac{x_2 + z}{y_2} \right) x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \\
& = y_1^{-1} \delta \left(\frac{x_1 + z}{y_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \tag{8.4}
\end{aligned}$$

and

$$\begin{aligned}
& x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) = \\
& = z^{-1} \delta \left(\frac{-x_1 + y_1}{z} \right) z^{-1} \delta \left(\frac{-x_2 + y_2}{z} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \\
& = \left(\sum_{m, n \in \mathbb{Z}} \frac{(-x_1 + y_1)^m}{z^{m+1}} \frac{(-x_2 + y_2)^n}{z^{n+1}} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \\
& = \left(\sum_{m, n \in \mathbb{Z}} (-x_2 + y_2)^{-1} \left(\frac{-x_1 + y_1}{-x_2 + y_2} \right)^m \frac{(-x_2 + y_2)^{m+n+1}}{z^{m+n+2}} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \\
& = \left(\sum_{m, k \in \mathbb{Z}} (-x_2 + y_2)^{-1} \left(\frac{-x_1 + y_1}{-x_2 + y_2} \right)^m z^{-1} \left(\frac{-x_2 + y_2}{z} \right)^k \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \\
& = (-x_2 + y_2)^{-1} \delta \left(\frac{-x_1 + y_1}{-x_2 + y_2} \right) z^{-1} \delta \left(\frac{-x_2 + y_2}{z} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \\
& = (-x_2)^{-1} \delta \left(\frac{x_1 - (y_1 - y_2)}{x_2} \right) z^{-1} \delta \left(\frac{-x_2 + y_2}{z} \right) y_1^{-1} \delta \left(\frac{y_2 + x_0}{y_1} \right) \\
& = x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) y_1^{-1} \delta \left(\frac{y_2 + x_0}{y_1} \right), \tag{8.5}
\end{aligned}$$

(8.3) becomes

$$\begin{aligned}
& ([Y'_{Q(z)}(v_1, x_1), Y'_{Q(z)}(v_2, x_2)] \lambda)(w_{(1)} \otimes w_{(2)}) = \\
& = \text{Res}_{x_0} \text{Res}_{y_2} \text{Res}_{y_1} y_1^{-1} \delta \left(\frac{x_1 + z}{y_1} \right) x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(Y(v_1, x_0)v_2, y_2)w_{(1)} \otimes w_{(2)}) \\
& \quad - \text{Res}_{x_0} \text{Res}_{y_2} \text{Res}_{y_1} x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) y_1^{-1} \delta \left(\frac{y_2 + x_0}{y_1} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(Y(v_1, x_0)v_2, y_2)w_{(2)}) \\
& = \text{Res}_{x_0} x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \cdot \\
& \quad \cdot \left(\text{Res}_{y_2} x_2^{-1} \delta \left(\frac{y_2 - z}{x_2} \right) \lambda(Y_1^*(Y(v_1, x_0)v_2, y_2)w_{(1)} \otimes w_{(2)}) \right. \\
& \quad \left. - \text{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \lambda(w_{(1)} \otimes Y_2(Y(v_1, x_0)v_2, y_2)w_{(2)}) \right)
\end{aligned}$$

$$= \text{Res}_{x_0} x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) (Y'_{Q(z)}(Y(v_1, x_0)v_2, x_2)\lambda)(w_{(1)} \otimes w_{(2)}). \quad (8.6)$$

Since λ , $w_{(1)}$ and $w_{(2)}$ are arbitrary, this equality gives the commutator formula for $Y'_{Q(z)}$, completing the proof of Proposition 5.2.

Remark 8.1 The proof of Proposition 5.2 suggests the following: Using the definitions (5.1) and (5.2) as motivation, let us define a (linear) action $\sigma_{Q(z)}$ of $V \otimes \iota_+ \mathbb{C}[t, t^{-1}, (z+t)^{-1}]$ on the vector space $W_1 \otimes W_2$ (as opposed to $(W_1 \otimes W_2)^*$) as follows:

$$\sigma_{Q(z)}(\xi)(w_{(1)} \otimes w_{(2)}) = \tau_{W_1}(T_{-z}^* \xi)w_{(1)} \otimes w_{(2)} - w_{(1)} \otimes \tau_{W_2}(T_{-z}^+ \xi)w_{(2)} \quad (8.7)$$

for $\xi \in V \otimes \iota_+ \mathbb{C}[t, t^{-1}, (z+t)^{-1}]$, $w_{(1)} \in W_1$, $w_{(2)} \in W_2$, or equivalently,

$$\begin{aligned} & \sigma_{Q(z)} \left(z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y_t(v, x_0) \right) (w_{(1)} \otimes w_{(2)}) \\ &= x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) Y_1^*(v, x_1)w_{(1)} \otimes w_{(2)} \\ & \quad - x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) w_{(1)} \otimes Y_2(v, x_1)w_{(2)}. \end{aligned} \quad (8.8)$$

That is, the operators $\sigma_{Q(z)}(\xi)$ and $\tau_{Q(z)}(\xi)$ are mutually adjoint:

$$(\tau_{Q(z)}(\xi)\lambda)(w_{(1)} \otimes w_{(2)}) = \lambda(\sigma_{Q(z)}(\xi)(w_{(1)} \otimes w_{(2)})). \quad (8.9)$$

While this action on $W_1 \otimes W_2$ is not very useful, it has the following three properties:

$$\sigma_{Q(z)}(Y_t(\mathbf{1}, x)) = 1, \quad (8.10)$$

$$\frac{d}{dx} \sigma_{Q(z)}(Y_t(v, x)) = \sigma_{Q(z)}(Y_t(L(-1)v, x)) \quad (8.11)$$

for $v \in V$,

$$\begin{aligned} & [\sigma_{Q(z)}(Y_t(v_2, x_2)), \sigma_{Q(z)}(Y_t(v_1, x_1))] \\ &= \text{Res}_{x_0} x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \sigma_{Q(z)}(Y_t(Y(v_1, x_0)v_2, x_2)) \end{aligned} \quad (8.12)$$

for $v_1, v_2 \in V$ (the *opposite commutator formula*). These follow either from the assertions of Propositions 5.1 and 5.2 or, better, from the fact that it was actually (8.10)–(8.12) that the proofs of these propositions were proving.

9 Some useful formulas

In this section, we shall prove some formulas which will be useful in the next section.

Let $\lambda \in (W_1 \otimes W_2)^*$, $w_{(1)} \in W_1$ and $w_{(2)} \in W_2$. From the definitions of $L'_{Q(z)}(0)$ and of $\tau_{Q(z)}$ (see Section 5), we have

$$\begin{aligned} (L'_{Q(z)}(0)\lambda)(w_{(1)} \otimes w_{(2)}) &= \\ &= (\tau_{Q(z)}(\omega \otimes t)\lambda)(w_{(1)} \otimes w_{(2)}) \\ &= \lambda((L(0) - zL(1))w_{(1)} \otimes w_{(2)}) - \lambda(w_{(1)} \otimes (L(0) - zL(-1))w_{(2)}), \end{aligned} \quad (9.1)$$

where we have used the same notations $L(0), L(-1), L(1)$ to denote the actions of the corresponding elements of the Virasoro algebra on both W_1 and W_2 . For convenience we write $L(-1) = L'_{Q(z)}(-1)$, $L(0) = L'_{Q(z)}(0)$ and $L(1) = L'_{Q(z)}(1)$ in the rest of this paper. There will be no confusion since the three $L(0)$'s act on different spaces.

Lemma 9.1 *For $\lambda \in (W_1 \otimes W_2)^*$, $w_{(1)} \in W_1$ and $w_{(2)} \in W_2$, we have*

$$\begin{aligned} \left(\left(1 - \frac{y_1}{z} \right)^{L(0)} \lambda \right) (w_{(1)} \otimes w_{(2)}) &= \\ &= \lambda \left(\left(1 - \frac{y_1}{z} \right)^{L(0)-zL(1)} w_{(1)} \otimes \left(1 - \frac{y_1}{z} \right)^{-(L(0)-zL(-1))} w_{(2)} \right). \end{aligned} \quad (9.2)$$

Proof From (9.1) we have

$$\begin{aligned} \left(\left(1 - \frac{y_1}{z} \right)^{L(0)} \lambda \right) (w_{(1)} \otimes w_{(2)}) &= \\ &= (e^{L(0) \log(1-\frac{y_1}{z})} \lambda) (w_{(1)} \otimes w_{(2)}) \\ &= \lambda(e^{L(0)-zL(1)} \log(1-\frac{y_1}{z}) w_{(1)} \otimes e^{-(L(0)-zL(-1)) \log(1-\frac{y_1}{z})} w_{(2)}) \\ &= \lambda \left(\left(1 - \frac{y_1}{z} \right)^{L(0)-zL(1)} w_{(1)} \otimes \left(1 - \frac{y_1}{z} \right)^{-(L(0)-zL(-1))} w_{(2)} \right). \quad \square \end{aligned} \quad (9.3)$$

Lemma 9.2 *For $v \in V$,*

$$Y'_{Q(z)}(v, x) = \left(1 - \frac{y_1}{z} \right)^{L(0)} Y'_{Q(z)} \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v, \frac{x}{1 - y_1/z} \right) \left(1 - \frac{y_1}{z} \right)^{-L(0)}. \quad (9.4)$$

This formula also holds for the vertex operators associated with any V -module.

Proof The identity (9.4) will follow from

$$e^{yL(0)}Y'_{Q(z)}(v, x)e^{-yL(0)} = Y'_{Q(z)}(e^{yL(0)}v, e^yx). \quad (9.5)$$

This is analogous to formula (5.2.37) of [FHL], but it is different because here we have e^y , not y ; also, the proof of (5.2.37) in [FHL] uses the grading of the module, which we do not have available here. To prove (9.5) assume without loss of generality that $\text{wt } v = h \in \mathbb{Z}$, and use the $L(-1)$ -derivative property (5.6) and the commutator formula (Proposition 5.2) to get

$$[L(0), Y'_{Q(z)}(v, x)] = \left(x \frac{d}{dx} + h\right) Y'_{Q(z)}(v, x), \quad (9.6)$$

which is the case $n = 0$ of (2.6.2) of [FHL]. Exactly as in Section 2.6 of [FHL], we obtain the desired result. $\square$

Lemma 9.3 *Let $L(-1)$, $L(0)$ be two operators satisfying the commutator relation*

$$[L(0), L(-1)] = L(-1). \quad (9.7)$$

Then

$$\left(1 - \frac{y_1}{z}\right)^{L(0)-zL(-1)} = e^{y_1L(-1)}\left(1 - \frac{y_1}{z}\right)^{L(0)}. \quad (9.8)$$

Proof We first prove that the derivative of

$$(1-x)^{L(0)-zL(-1)}(1-x)^{-L(0)}e^{-zxL(-1)}$$

is 0. Write $A = (1-x)^{L(0)-zL(-1)}$, $B = (1-x)^{-L(0)}$, $C = e^{-zxL(-1)}$. Then

$$\begin{aligned} \frac{d}{dx}(ABC) &= -A(1-x)^{-1}(L(0) - zL(-1))BC \\ &\quad + A(1-x)^{-1}L(0)BC \\ &\quad - zABL(-1)C. \end{aligned} \quad (9.9)$$

Using (5.2.12) of [FHL] we have

$$\begin{aligned} BL(-1) &= (1-x)^{-L(0)}L(-1) \\ &= e^{(-\log(1-x))L(0)}L(-1) \\ &= L(-1)e^{(-\log(1-x))L(0)}e^{-\log(1-x)} \\ &= L(-1)(1-x)^{-L(0)}(1-x)^{-1} \\ &= (1-x)^{-1}L(-1)B. \end{aligned} \quad (9.10)$$

Substituting (9.10) into (9.9), we obtain

$$\begin{aligned}\frac{d}{dx}(ABC) &= -A(1-x)^{-1}L(0)BC + zA(1-x)^{-1}L(-1)BC \\ &\quad + A(1-x)^{-1}L(0)BC - zA(1-x)^{-1}L(-1)BC \\ &= 0.\end{aligned}$$

Thus ABC is constant, and since $ABC\Big|_{x=0} = 1$, we have $ABC = 1$, which is equivalent to (9.8). $\square$

10 Jacobi identity for vertex operators on a certain subspace of $(W_1 \otimes W_2)^*$

We prove Theorem 6.1 in this section. The reader should again observe the justifiability of each formal step in the argument; sometimes this is quite subtle.

Let λ be an element of $(W_1 \otimes W_2)^*$ satisfying the compatibility condition, that is, (a) the lower truncation condition: for all $v \in V$,

$$\tau_{Q(z)}(v \otimes t^n)\lambda = 0 \quad \text{for } n \text{ sufficiently large,} \quad (10.1)$$

or equivalently, for all $v \in V$,

$$\begin{aligned}Y'_{Q(z)}(v, x)\lambda &= \tau_{Q(z)}(Y_t(v, x))\lambda \\ &= \sum_{n < N} \tau_{Q(z)}(v \otimes t^n)\lambda x^{-n-1} \quad \text{for some } N \in \mathbb{Z},\end{aligned} \quad (10.2)$$

and (b) the formula

$$\begin{aligned}\tau_{Q(z)}\left(x_1^{-1}\delta\left(\frac{z+x_0}{x_1}\right)Y_t(v, x_0)\right)\lambda &= \\ &= x_1^{-1}\delta\left(\frac{z+x_0}{x_1}\right)Y'_{Q(z)}(v, x_0)\lambda \quad \text{for all } v \in V.\end{aligned} \quad (10.3)$$

By (5.2) and (5.4), (10.3) is equivalent to

$$x_0^{-1}\delta\left(\frac{x_1-z}{x_0}\right)\lambda(Y_1^*(v, x_1)w_{(1)} \otimes w_{(2)})$$

$$\begin{aligned}
& -x_0^{-1}\delta\left(\frac{z-x_1}{-x_0}\right)\lambda(w_{(1)}\otimes Y_2(v,x_1)w_{(2)}) \\
& = x_1^{-1}\delta\left(\frac{z+x_0}{x_1}\right)\left(\operatorname{Res}_{y_1}x_0^{-1}\delta\left(\frac{y_1-z}{x_0}\right)\lambda(Y_1^*(v,y_1)w_{(1)}\otimes w_{(2)})\right. \\
& \quad \left.-\operatorname{Res}_{y_1}x_0^{-1}\delta\left(\frac{z-y_1}{-x_0}\right)\lambda(w_{(1)}\otimes Y_2(v,y_1)w_{(2)})\right) \tag{10.4}
\end{aligned}$$

for all $v \in V$, $w_{(1)} \in W_1$ and $w_{(2)} \in W_2$. Note that on the right-hand side the distributive law is not valid since the two individual products are not defined. One feature of the argument which follows is that we must rewrite expressions to allow the application of distributivity.

By (5.4), we have

$$\begin{aligned}
& \left(x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)Y'_{Q(z)}(v_1,x_1)Y'_{Q(z)}(v_2,x_2)\lambda\right)(w_{(1)}\otimes w_{(2)}) \\
& = x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)(Y'_{Q(z)}(v_1,x_1)Y'_{Q(z)}(v_2,x_2)\lambda)(w_{(1)}\otimes w_{(2)}) \\
& = x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)\left(\operatorname{Res}_{y_1}x_1^{-1}\delta\left(\frac{y_1-z}{x_1}\right)\right. \\
& \quad \cdot (Y'_{Q(z)}(v_2,x_2)\lambda)(Y_1^*(v_1,y_1)w_{(1)}\otimes w_{(2)}) \\
& \quad \left.-\operatorname{Res}_{y_1}x_1^{-1}\delta\left(\frac{z-y_1}{-x_1}\right)(Y'_{Q(z)}(v_2,x_2)\lambda)(w_{(1)}\otimes Y_2(v_1,y_1)w_{(2)})\right) \\
& = x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)\left(\operatorname{Res}_{y_1}x_1^{-1}\delta\left(\frac{y_1-z}{x_1}\right)\operatorname{Res}_{y_2}x_2^{-1}\delta\left(\frac{y_2-z}{x_2}\right)\right. \\
& \quad \cdot \lambda(Y_1^*(v_2,y_2)Y_1^*(v_1,y_1)w_{(1)}\otimes w_{(2)}) \\
& \quad \left.-\operatorname{Res}_{y_1}x_1^{-1}\delta\left(\frac{y_1-z}{x_1}\right)\operatorname{Res}_{y_2}x_2^{-1}\delta\left(\frac{z-y_2}{-x_2}\right)\right. \\
& \quad \cdot \lambda(Y_1^*(v_1,y_1)w_{(1)}\otimes Y_2(v_2,y_2)w_{(2)}) \\
& \quad \left.-\operatorname{Res}_{y_1}x_1^{-1}\delta\left(\frac{z-y_1}{-x_1}\right)(Y'_{Q(z)}(v_2,x_2)\lambda)(w_{(1)}\otimes Y_2(v_1,y_1)w_{(2)})\right). \tag{10.5}
\end{aligned}$$

From the properties of the formal δ -function, we see that the right-hand side of (10.5) is equal to

$$\begin{aligned}
& x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)\left(\operatorname{Res}_{y_1}y_1^{-1}\delta\left(\frac{x_1+z}{y_1}\right)\operatorname{Res}_{y_2}y_2^{-1}\delta\left(\frac{x_2+z}{y_2}\right)\right. \\
& \quad \left.\cdot \lambda(Y_1^*(v_2,x_2+z)Y_1^*(v_1,x_1+z)w_{(1)}\otimes w_{(2)})\right)
\end{aligned}$$

$$\begin{aligned}
& - \operatorname{Res}_{y_1} y_1^{-1} \delta \left(\frac{x_1 + z}{y_1} \right) \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_1, x_1 + z)w_{(1)} \otimes Y_2(v_2, y_2)w_{(2)}) \\
& - \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) (Y'_{Q(z)}(v_2, x_2)\lambda)(w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& = x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \left(\lambda(Y_1^*(v_2, x_2 + z)Y_1^*(v_1, x_1 + z)w_{(1)} \otimes w_{(2)}) \right. \\
& \quad - \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \lambda(Y_1^*(v_1, x_1 + z)w_{(1)} \otimes Y_2(v_2, y_2)w_{(2)}) \\
& \quad \left. - \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) (Y'_{Q(z)}(v_2, x_2)\lambda)(w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \right). \quad (10.6)
\end{aligned}$$

From the $L(-1)$ -derivative property for Y_1 and Y_1^* and the commutator formulas for $L(-1)$, $Y_1(\cdot, x)$ and for $L(1)$, $Y_1^*(\cdot, x)$, we obtain

$$\begin{aligned}
Y_1(v, x + z) &= Y_1(e^{zL(-1)}v, x) \\
&= e^{zL(-1)}Y_1(v, x)e^{-zL(-1)} \\
&= \sum_{n \geq 0} \frac{z^n}{n!} \frac{d^n}{dx^n} Y_1(v, x), \quad (10.7)
\end{aligned}$$

and

$$\begin{aligned}
Y_1^*(v, x + z) &= Y_1^*(e^{zL(-1)}v, x) \\
&= e^{-zL(1)}Y_1^*(v, x)e^{zL(1)} \\
&= \sum_{n \geq 0} \frac{z^n}{n!} \frac{d^n}{dx^n} Y_1^*(v, x). \quad (10.8)
\end{aligned}$$

(Note that all these expressions are in fact defined.) Using (10.8), we see that the right-hand side of (10.6) can be written as

$$\begin{aligned}
& x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \left(\lambda(Y_1^*(e^{zL(-1)}v_2, x_2)Y_1^*(e^{zL(-1)}v_1, x_1)w_{(1)} \otimes w_{(2)}) \right. \\
& \quad - \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \lambda(Y_1^*(e^{zL(-1)}v_1, x_1)w_{(1)} \otimes Y_2(v_2, y_2)w_{(2)}) \\
& \quad \left. - \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) (Y'_{Q(z)}(v_2, x_2)\lambda)(w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \right) \\
& = x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \left(\lambda(e^{-zL(1)}Y_1^*(v_2, x_2)Y_1^*(v_1, x_1)e^{zL(1)}w_{(1)} \otimes w_{(2)}) \right)
\end{aligned}$$

$$\begin{aligned}
& - \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \lambda(e^{-zL(1)} Y_1^*(v_1, x_1) e^{zL(1)} w_{(1)} \otimes Y_2(v_2, y_2) w_{(2)}) \\
& - \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) (Y'_{Q(z)}(v_2, x_2) \lambda)(w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}). \quad (10.9)
\end{aligned}$$

Note that it is easier to verify the well-definedness of the terms on the right-hand side of (10.6) than that of the terms in (10.9), though (10.9) is sometimes easier to use if it is known that every term is well defined. Below we shall write expressions like those on the right-hand side of (10.6) in whichever way suits our needs. The distributive law applies to the right-hand side of (10.6) (or (10.9)) since all three of the following expressions are defined:

$$\begin{aligned}
& x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \lambda(Y_1^*(v_2, x_2 + z) Y_1^*(v_1, x_1 + z) w_{(1)} \otimes w_{(2)}), \\
& x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \lambda(Y_1^*(v_1, x_1 + z) w_{(1)} \otimes Y_2(v_2, y_2) w_{(2)}), \\
& x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) (Y'_{Q(z)}(v_2, x_2) \lambda)(w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}).
\end{aligned}$$

Now we examine the last expression in (10.9). Rewriting the formal δ -functions $x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right)$ and $x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right)$, and using (9.4) and the fundamental property of the δ -function, we have:

$$\begin{aligned}
& x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot (Y'_{Q(z)}(v_2, x_2) \lambda)(w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
& = \left(\frac{x_1}{z} \right)^{-1} \left(\frac{x_0}{x_1/z} \right)^{-1} \delta \left(\frac{z + \left(\frac{x_2}{-x_1/z} \right)}{\frac{x_0}{x_1/z}} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) \cdot \\
& \quad \cdot \left(\left(1 - \frac{y_1}{z} \right)^{L(0)} Y'_{Q(z)} \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v_2, \frac{x_2}{1 - y_1/z} \right) \cdot \right. \\
& \quad \cdot \left. \left(1 - \frac{y_1}{z} \right)^{-L(0)} \lambda \right) (w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
& = \left(\frac{x_1}{z} \right)^{-1} \left(\frac{x_0}{x_1/z} \right)^{-1} \delta \left(\frac{z + \left(\frac{x_2}{-x_1/z} \right)}{\frac{x_0}{x_1/z}} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) \cdot \\
& \quad \cdot \left(\left(1 - \frac{y_1}{z} \right)^{L(0)} Y'_{Q(z)} \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v_2, \frac{x_2}{-x_1/z} \right) \cdot \right.
\end{aligned}$$

$$\cdot \left(1 - \frac{y_1}{z}\right)^{-L(0)} \lambda \Big) (w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}). \quad (10.10)$$

By (9.2) and (10.3), the right-hand side of (10.10) is equal to

$$\begin{aligned} & \left(\frac{x_1}{z}\right)^{-1} \left(\frac{x_0}{x_1/z}\right)^{-1} \delta \left(\frac{z + \left(\frac{x_2}{-x_1/z}\right)}{\frac{x_0}{x_1/z}} \right) \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) \cdot \\ & \cdot \left(Y'_{Q(z)} \left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} v_2, \frac{x_2}{-x_1/z} \right) \left(1 - \frac{y_1}{z}\right)^{-L(0)} \lambda \right) \cdot \\ & \cdot \left(\left(1 - \frac{y_1}{z}\right)^{L(0)-zL(1)} w_{(1)} \otimes \right. \\ & \quad \left. \otimes \left(1 - \frac{y_1}{z}\right)^{-L(0)-zL(-1)} Y_2(v_1, y_1) w_{(2)} \right) \\ & = \left(\frac{x_1}{z}\right)^{-1} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) \cdot \\ & \cdot \left(\tau_{Q(z)} \left(\left(\frac{x_0}{x_1/z}\right)^{-1} \delta \left(\frac{z + \left(\frac{x_2}{-x_1/z}\right)}{\frac{x_0}{x_1/z}} \right) \cdot \right. \right. \\ & \quad \left. \cdot Y_t \left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} v_2, \frac{x_2}{-x_1/z} \right) \right) \left(1 - \frac{y_1}{z}\right)^{-L(0)} \lambda \Big) \cdot \\ & \cdot \left(\left(1 - \frac{y_1}{z}\right)^{L(0)-zL(1)} w_{(1)} \otimes \right. \\ & \quad \left. \otimes \left(1 - \frac{y_1}{z}\right)^{-L(0)-zL(-1)} Y_2(v_1, y_1) w_{(2)} \right). \quad (10.11) \end{aligned}$$

By (5.2), the right-hand side of (10.11) becomes

$$\begin{aligned} & \left(\frac{x_1}{z}\right)^{-1} \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) \left(\left(\frac{x_2}{-x_1/z}\right)^{-1} \delta \left(\frac{\frac{x_0}{x_1/z} - z}{\frac{x_2}{-x_1/z}} \right) \cdot \right. \\ & \cdot \left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} \lambda \right) \left(Y_1^* \left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} v_2, \frac{x_0}{x_1/z} \right) \cdot \right. \\ & \quad \left. \cdot \left(1 - \frac{y_1}{z}\right)^{L(0)-zL(1)} w_{(1)} \otimes \left(1 - \frac{y_1}{z}\right)^{-(L(0)-zL(1))} Y_2(v_1, y_1) w_{(2)} \right) \\ & - \left(\frac{x_2}{-x_1/z} \right)^{-1} \delta \left(\frac{z - \frac{x_0}{x_1/z}}{\frac{x_2}{-x_1/z}} \right) \left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} \lambda \right) \left(\left(1 - \frac{y_1}{z}\right)^{L(0)-zL(1)} \cdot \right. \end{aligned}$$

$$\begin{aligned}
& \cdot w_{(1)} \otimes Y_2 \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v_2, \frac{x_0}{x_1/z} \right) \cdot \\
& \cdot \left(1 - \frac{y_1}{z} \right)^{(L(0)-zL(-1))} Y_2(v_1, y_1) w_{(2)} \Big). \tag{10.12}
\end{aligned}$$

Using (9.2) again but with $1 - \frac{y_1}{z}$ replaced by $(1 - \frac{y_1}{z})^{-1}$, rewriting formal δ -functions and then using the distributive law, we see that (10.12) is equal to

$$\begin{aligned}
& \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) \left(-x_2^{-1} \delta \left(\frac{x_0 - x_1}{-x_2} \right) \cdot \right. \\
& \quad \cdot \lambda \left(\left(1 - \frac{y_1}{z} \right)^{-(L(0)-zL(1))} Y_1^* \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v_2, \frac{x_0}{-(1 - y_1/z)} \right) \cdot \right. \\
& \quad \cdot \left(1 - \frac{y_1}{z} \right)^{L(0)-zL(1)} w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)} \Big) \\
& \quad + x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \lambda(w_{(1)} \otimes \left(1 - \frac{y_1}{z} \right)^{L(0)-zL(-1)} \cdot \\
& \quad \cdot Y_2 \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v_2, \frac{x_0}{-(1 - y_1/z)} \right) \cdot \\
& \quad \cdot \left(1 - \frac{y_1}{z} \right)^{-(L(0)-zL(-1))} Y_2(v_1, y_1) w_{(2)} \Big) \\
& = - \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) x_2^{-1} \delta \left(\frac{x_0 - x_1}{-x_2} \right) \cdot \\
& \quad \cdot \lambda \left(\left(1 - \frac{y_1}{z} \right)^{-(L(0)-zL(1))} Y_1^* \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v_2, \frac{x_0}{-(1 - y_1/z)} \right) \cdot \right. \\
& \quad \cdot \left(1 - \frac{y_1}{z} \right)^{L(0)-zL(1)} w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)} \Big) \\
& \quad + \text{Res}_{y_1} x_1^{-1} \delta \left(\frac{1 - y_1/z}{-x_1/z} \right) x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \lambda(w_{(1)} \otimes \\
& \quad \otimes \left(1 - \frac{y_1}{z} \right)^{L(0)-zL(-1)} Y_2 \left(\left(1 - \frac{y_1}{z} \right)^{-L(0)} v_2, \frac{x_0}{-(1 - y_1/z)} \right) \cdot \\
& \quad \cdot \left(1 - \frac{y_1}{z} \right)^{-(L(0)-zL(-1))} Y_2(v_1, y_1) w_{(2)} \Big). \tag{10.13}
\end{aligned}$$

But by (9.8) and (9.4),

$$\begin{aligned}
& \left(1 - \frac{y_1}{z}\right)^{L(0)-zL(-1)} Y_2\left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} v_2, \frac{x_0}{-(1-y_1/z)}\right) \left(1 - \frac{y_1}{z}\right)^{-(L(0)-zL(-1))} \\
&= e^{y_1 L(-1)} \left(1 - \frac{y_1}{z}\right)^{L(0)} Y_2\left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} v_2, \frac{x_0}{-(1-y_1/z)}\right) \cdot \\
&\quad \cdot \left(1 - \frac{y_1}{z}\right)^{-L(0)} e^{-y_1 L(-1)} \\
&= e^{y_1 L(-1)} Y_2(v_2, -x_0) e^{-y_1 L(-1)} \\
&= Y_2(v_2, -x_0 + y_1) \\
&= Y_2(v_2, -x_0 - (z - y_1) + z) \\
&= Y_2(e^{zL(-1)} v_2, -x_0 - (z - y_1)). \tag{10.14}
\end{aligned}$$

We similarly have (using (9.8) and (9.4) for $Y_1'(v_2, x)$ and then using (3.22))

$$\begin{aligned}
& \left(1 - \frac{y_1}{z}\right)^{-(L(0)-zL(1))} Y_1^*\left(\left(1 - \frac{y_1}{z}\right)^{-L(0)} v_2, \frac{x_0}{-(1-y_1/z)}\right) \left(1 - \frac{y_1}{z}\right)^{L(0)-zL(1)} \\
&= e^{-y_1 L(1)} Y_1^*(v_2, -x_0) e^{y_1 L(1)} \\
&= Y_1^*(v_2, -x_0 + y_1) \\
&= Y_1^*(v_2, -x_0 - (z - y_1) + z) \\
&= Y_1^*(e^{zL(-1)} v_2, -x_0 - (z - y_1)). \tag{10.15}
\end{aligned}$$

Substituting (10.14) and (10.15) into the right-hand side of (10.13) and then combining with (10.10)–(10.13), we obtain

$$\begin{aligned}
& x_0^{-1} \delta\left(\frac{x_1 - x_2}{x_0}\right) \text{Res}_{y_1} x_1^{-1} \delta\left(\frac{z - y_1}{-x_1}\right) (Y'_{Q(z)}(v_2, x_2) \lambda)(w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
&= - \text{Res}_{y_1} x_1^{-1} \delta\left(\frac{z - y_1}{-x_1}\right) x_2^{-1} \delta\left(\frac{x_0 - x_1}{-x_2}\right) \cdot \\
&\quad \cdot \lambda(Y_1^*\left(v_2, -x_0 - (z - y_1) + z\right) w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
&\quad + \text{Res}_{y_1} x_1^{-1} \delta\left(\frac{z - y_1}{-x_1}\right) x_2^{-1} \delta\left(\frac{x_1 - x_0}{x_2}\right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, -x_0 + y_1) Y_2(v_1, y_1) w_{(2)}). \tag{10.16}
\end{aligned}$$

(We choose the form of the expression from (10.15) in anticipation of the next step.)

By the properties of the formal δ -function, the right-hand side of (10.16) is equal to

$$\begin{aligned}
& -x_2^{-1} \delta \left(\frac{-x_0 + x_1}{x_2} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(v_2, -x_0 + x_1 + z)w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& + x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, -x_0 + y_1)Y_2(v_1, y_1)w_{(2)}) \\
& = -x_2^{-1} \delta \left(\frac{-x_0 + x_1}{x_2} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(e^{zL(-1)}v_2, x_2)w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& + x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, -x_0 + y_1)Y_2(v_1, y_1)w_{(2)}). \tag{10.17}
\end{aligned}$$

Since

$$\operatorname{Res}_{y_2} y_2^{-1} \delta \left(\frac{-x_0 + y_1}{y_2} \right) = 1,$$

the right-hand side of (10.17) can be written as

$$\begin{aligned}
& -x_2^{-1} \delta \left(\frac{-x_0 + x_1}{x_2} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(e^{zL(-1)}v_2, x_2)w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& + x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \operatorname{Res}_{y_2} y_2^{-1} \delta \left(\frac{-x_0 + y_1}{y_2} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, -x_0 + y_1)Y_2(v_1, y_1)w_{(2)}). \tag{10.18}
\end{aligned}$$

Again by the properties of the formal δ -function, (10.18) becomes

$$\begin{aligned}
& x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(e^{zL(-1)}v_2, x_2)w_{(1)} \otimes Y_2(v_1, y_1)w_{(2)}) \\
& + x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \operatorname{Res}_{y_2} y_2^{-1} \delta \left(\frac{-x_0 + y_1}{y_2} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, y_2)Y_2(v_1, y_1)w_{(2)})
\end{aligned}$$

$$\begin{aligned}
&= x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
&\quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_2, x_2) w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
&\quad + x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} \operatorname{Res}_{y_2} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) y_2^{-1} \delta \left(\frac{-x_0 + y_1}{y_2} \right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, y_2) Y_2(v_1, y_1) w_{(2)}) \\
&= x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
&\quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_2, x_2) w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
&\quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} \operatorname{Res}_{y_2} \cdot \\
&\quad \cdot (x_2 + x_0)^{-1} \delta \left(\frac{z - y_1}{-x_2 - x_0} \right) x_0^{-1} \delta \left(\frac{y_2 - y_1}{-x_0} \right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, y_2) Y_2(v_1, y_1) w_{(2)}) \\
&= x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
&\quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_2, x_2) w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
&\quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) x_0^{-1} \delta \left(\frac{y_2 - y_1}{-x_0} \right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, y_2) Y_2(v_1, y_1) w_{(2)}). \tag{10.19}
\end{aligned}$$

Substituting (10.16)–(10.19) into (10.9) we obtain

$$\begin{aligned}
&\left(x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y'_{Q(z)}(v_1, x_1) Y'_{Q(z)}(v_2, x_2) \lambda \right) (w_{(1)} \otimes w_{(2)}) \\
&= x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \lambda(Y_1^*(e^{zL(-1)} v_2, x_2) Y_1^*(e^{zL(-1)} v_1, x_1) w_{(1)} \otimes w_{(2)}) \\
&\quad - x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
&\quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_1, x_1) w_{(1)} \otimes Y_2(v_2, x_2) w_{(2)}) \\
&\quad - x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
&\quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_2, x_2) w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
&\quad + x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) x_0^{-1} \delta \left(\frac{y_2 - y_1}{-x_0} \right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(v_2, y_2) Y_2(v_1, y_1) w_{(2)}). \tag{10.20}
\end{aligned}$$

Now consider the result of the calculation from (10.16) to (10.19) except for the last two steps in (10.19). Reversing the subscripts 1 and 2 of the symbols v , x and y and replacing x_0 by $-x_0$ in this result and then using a property of the formal δ -function, we have

$$\begin{aligned}
& x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot (Y'_{Q(z)}(v_1, x_1) \lambda)(w_{(1)} \otimes Y_2(v_2, y_2) w_{(2)}) \\
& = x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_1, x_1) w_{(1)} \otimes Y_2(v_2, x_2) w_{(2)}) \\
& \quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) x_0^{-1} \delta \left(\frac{y_1 - y_2}{x_0} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_1, y_1) Y_2(v_2, y_2) w_{(2)}). \tag{10.21}
\end{aligned}$$

From (10.5)–(10.9), again reversing the subscripts 1 and 2 of the symbols v , x and y and replacing x_0 by $-x_0$, and (10.21), we have

$$\begin{aligned}
& \left(-x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) Y'_{Q(z)}(v_2, x_2) Y'_{Q(z)}(v_1, x_1) \cdot \lambda \right) (w_{(1)} \otimes w_{(2)}) \\
& = -x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \lambda(Y_1^*(e^{zL(-1)} v_1, x_1) Y_1^*(e^{zL(-1)} v_2, x_2) w_{(1)} \otimes w_{(2)}) \\
& \quad + x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \operatorname{Res}_{y_1} x_1^{-1} \delta \left(\frac{z - y_1}{-x_1} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_2, x_2) w_{(1)} \otimes Y_2(v_1, y_1) w_{(2)}) \\
& \quad + x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
& \quad \cdot \lambda(Y_1^*(e^{zL(-1)} v_1, x_1) w_{(1)} \otimes Y_2(v_2, x_2) w_{(2)}) \\
& \quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \operatorname{Res}_{y_1} \operatorname{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) x_0^{-1} \delta \left(\frac{y_1 - y_2}{x_0} \right) \cdot \\
& \quad \cdot \lambda(w_{(1)} \otimes Y_2(v_1, y_1) Y_2(v_2, y_2) w_{(2)}). \tag{10.22}
\end{aligned}$$

The formulas (10.20) and (10.22) give:

$$\begin{aligned}
& \left(\left(x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y'_{Q(z)}(v_1, x_1) Y'_{Q(z)}(v_2, x_2) \right. \right. \\
& \quad \left. \left. - x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) Y'_{Q(z)}(v_2, x_2) Y'_{Q(z)}(v_1, x_1) \right) \lambda \right) (w_{(1)} \otimes w_{(2)})
\end{aligned}$$

$$\begin{aligned}
&= \lambda \left(\left(x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y_1^*(e^{zL(-1)} v_2, x_2) Y_1^*(e^{zL(-1)} v_1, x_1) \right. \right. \\
&\quad \left. \left. - x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) Y_1^*(e^{zL(-1)} v_1, x_1) Y_1^*(e^{zL(-1)} v_2, x_2) \right) w_{(1)} \otimes w_{(2)} \right) \\
&\quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \text{Res}_{y_1} \text{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
&\quad \cdot \lambda \left(w_{(1)} \otimes \left(x_0^{-1} \delta \left(\frac{y_1 - y_2}{x_0} \right) Y_2(v_1, y_1) Y_2(v_2, y_2) \right. \right. \\
&\quad \left. \left. - x_0^{-1} \delta \left(\frac{y_2 - y_1}{-x_0} \right) Y_2(v_2, y_2) Y_2(v_1, y_1) \right) w_{(2)} \right). \tag{10.23}
\end{aligned}$$

From the Jacobi identities for Y_1^* and Y_2 and the exponentiated form of the formula for the commutator of $L(-1)$ and the vertex operators $Y(\cdot, x_0)$ (cf. (10.7)), the right-hand side of (10.23) is equal to

$$\begin{aligned}
&x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \lambda(Y_1^*(Y(e^{zL(-1)} v_1, x_0) e^{zL(-1)} v_2, x_2) w_{(1)} \otimes w_{(2)}) \\
&\quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \text{Res}_{y_1} \text{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(Y(v_1, x_0) v_2, y_2) w_{(2)}) \\
&= x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \lambda(Y_1^*(e^{zL(-1)} Y(v_1, x_0) v_2, x_2) w_{(1)} \otimes w_{(2)}) \\
&\quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \text{Res}_{y_1} \text{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) y_2^{-1} \delta \left(\frac{y_1 - x_0}{y_2} \right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(Y(v_1, x_0) v_2, y_2) w_{(2)}). \tag{10.24}
\end{aligned}$$

Using (10.8), evaluating Res_{y_1} and then using the definition of $Y'_{Q(z)}$ (recall (5.4)), we finally see that the right-hand side of (10.24) is equal to

$$\begin{aligned}
&x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \lambda(Y_1^*(Y(v_1, x_0) v_2, x_2 + z) w_{(1)} \otimes w_{(2)}) \\
&\quad - x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) \text{Res}_{y_2} x_2^{-1} \delta \left(\frac{z - y_2}{-x_2} \right) \cdot \\
&\quad \cdot \lambda(w_{(1)} \otimes Y_2(Y(v_1, x_0) v_2, y_2) w_{(2)}) \\
&= x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) (Y'_{Q(z)}(Y(v_1, x_0) v_2, x_2) \lambda)(w_{(1)} \otimes w_{(2)}), \tag{10.25}
\end{aligned}$$

proving Theorem 6.1 in [HL].

11 Stability of certain subspaces of $(W_1 \otimes W_2)^*$ under the action of vertex operators

We prove Proposition 6.2 in this section.

Let λ be an element of $(W_1 \otimes W_2)^*$ satisfying the compatibility condition. We first want to prove that each coefficient in x of $Y'_{Q(z)}(u, x_0)Y'_{Q(z)}(v, x)\lambda$ is a Laurent series involving only finitely many negative powers of x_0 and that

$$\begin{aligned} \tau_{Q(z)} \left(z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y_t(u, x_0) \right) Y'_{Q(z)}(v, x)\lambda \\ = z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y'_{Q(z)}(u, x_0)Y'_{Q(z)}(v, x)\lambda \end{aligned} \quad (11.1)$$

for all $u, v \in V$. Using the commutator formula for $Y'_{Q(z)}$, we have

$$\begin{aligned} Y'_{Q(z)}(u, x_0)Y'_{Q(z)}(v, x)\lambda &= \\ &= Y'_{Q(z)}(v, x)Y'_{Q(z)}(u, x_0)\lambda \\ &\quad - \text{Res}_y x_0^{-1} \delta \left(\frac{x - y}{x_0} \right) Y'_{Q(z)}(Y(v, y)u, x_0)\lambda. \end{aligned} \quad (11.2)$$

Each coefficient in x of the right-hand side of (11.2) is a Laurent series involving only finitely many negative powers of x_0 since λ satisfies the lower truncation condition. Thus the coefficients in x of $Y'_{Q(z)}(v, x)\lambda$ satisfy the lower truncation condition.

By (5.2) and (5.4), we have

$$\begin{aligned} &\left(\tau_{Q(z)} \left(z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y_t(u, x_0) \right) Y'_{Q(z)}(v, x)\lambda \right) (w_{(1)} \otimes w_{(2)}) \\ &= x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) (Y'_{Q(z)}(v, x)\lambda) (Y_1^*(u, x_1)w_{(1)} \otimes w_{(2)}) \\ &\quad - x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) (Y'_{Q(z)}(v, x)\lambda) (w_{(1)} \otimes Y_2(u, x_1)w_{(2)}) \\ &= x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \left(\lambda(Y_1^*(v, x + z)Y_1^*(u, x_1)w_{(1)} \otimes w_{(2)}) \right. \\ &\quad \left. - \text{Res}_{x_2} x^{-1} \delta \left(\frac{z - x_2}{-x} \right) \lambda(Y_1^*(u, x_1)w_{(1)} \otimes Y_2(v, x_2)w_{(2)}) \right) \\ &\quad - x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \left(\lambda(Y_1^*(e^{zL(-1)}v, x)w_{(1)} \otimes Y_2(u, x_1)w_{(2)}) \right. \\ &\quad \left. - \text{Res}_{x_2} x^{-1} \delta \left(\frac{z - x_2}{-x} \right) \lambda(w_{(1)} \otimes Y_2(v, x_2)Y_2(u, x_1)w_{(2)}) \right). \end{aligned} \quad (11.3)$$

Now the distributive law applies, giving us four terms. Inserting

$$\text{Res}_{x_4} x_4^{-1} \delta \left(\frac{x+z}{x_4} \right) = 1$$

into the first of these terms and correspondingly replacing $x+z$ by x_4 in $Y_1^*(v, x+z)$, we can apply the commutator formula for Y_1^* in the usual way. Also using the commutator formula for Y_2 and properties of the δ -function, we write the right-hand side of (11.3) as

$$\begin{aligned} & x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \lambda(Y_1^*(u, x_1) Y_1^*(v, x+z) w_{(1)} \otimes w_{(2)}) \\ & - x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \text{Res}_{x_4} \text{Res}_{x_3} x_1^{-1} \delta \left(\frac{x_4 - x_3}{x_1} \right) x_4^{-1} \delta \left(\frac{x+z}{x_4} \right) \cdot \\ & \quad \cdot \lambda(Y_1^*(Y(v, x_3)u, x_1) w_{(1)} \otimes w_{(2)}) \\ & - x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \text{Res}_{x_2} x^{-1} \delta \left(\frac{z - x_2}{-x} \right) \lambda(Y_1^*(u, x_1) w_{(1)} \otimes Y_2(v, x_2) w_{(2)}) \\ & - x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \lambda(Y_1^*(e^{zL(-1)}v, x) w_{(1)} \otimes Y_2(u, x_1) w_{(2)}) \\ & + x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \text{Res}_{x_2} x^{-1} \delta \left(\frac{z - x_2}{-x} \right) \lambda(w_{(1)} \otimes Y_2(u, x_1) Y_2(v, x_2) w_{(2)}) \\ & + x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \text{Res}_{x_2} x^{-1} \delta \left(\frac{z - x_2}{-x} \right) \text{Res}_{x_3} x_1^{-1} \delta \left(\frac{x_2 - x_3}{x_1} \right) \cdot \\ & \quad \cdot \lambda(w_{(1)} \otimes Y_2(Y(v, x_3)u, x_1) w_{(2)}) \\ & = \left(x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \lambda(Y_1^*(u, x_1) Y_1^*(e^{zL(-1)}v, x) w_{(1)} \otimes w_{(2)}) \right. \\ & \quad \left. - x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \lambda(Y_1^*(e^{zL(-1)}v, x) w_{(1)} \otimes Y_2(u, x_1) w_{(2)}) \right) \\ & - \text{Res}_{x_2} x^{-1} \delta \left(\frac{z - x_2}{-x} \right) \left(x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \cdot \right. \\ & \quad \cdot \lambda(Y_1^*(u, x_1) w_{(1)} \otimes Y_2(v, x_2) w_{(2)}) \\ & \quad \left. - x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \lambda(w_{(1)} \otimes Y_2(u, x_1) Y_2(v, x_2) w_{(2)}) \right) \\ & - \left(\text{Res}_{x_4} \text{Res}_{x_3} (x_0 + z)^{-1} \delta \left(\frac{x + z - x_3}{x_0 + z} \right) \cdot \right. \\ & \quad \cdot x_4^{-1} \delta \left(\frac{x + z}{x_4} \right) x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \cdot \\ & \quad \cdot \lambda(Y_1^*(Y(v, x_3)u, x_1) w_{(1)} \otimes w_{(2)}) \end{aligned}$$

$$\begin{aligned}
& - \operatorname{Res}_{x_2} \operatorname{Res}_{x_3} x^{-1} \delta \left(\frac{z - (x_1 + x_3)}{-x} \right) x_2^{-1} \delta \left(\frac{x_1 + x_3}{x_2} \right) \cdot \\
& \cdot x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \lambda(w_{(1)} \otimes Y_2(Y(v, x_3)u, x_1)w_{(2)}). \quad (11.4)
\end{aligned}$$

By (5.2) and properties of formal δ -function, the right-hand side of (11.4) is equal to

$$\begin{aligned}
& \left(\tau_{Q(z)} \left(z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y_t(u, x_0) \right) \lambda \right) (Y_1^*(e^{zL(-1)}v, x)w_{(1)} \otimes w_{(2)}) \\
& - \operatorname{Res}_{x_2} x^{-1} \delta \left(\frac{z - x_2}{-x} \right) \cdot \\
& \cdot \left(\tau_{Q(z)} \left(z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y_t(u, x_0) \right) \lambda \right) (w_{(1)} \otimes Y_2(v, x_2)w_{(2)}) \\
& - \left(\operatorname{Res}_{x_3} x_0^{-1} \delta \left(\frac{x - x_3}{x_0} \right) \cdot \right. \\
& \cdot x_0^{-1} \delta \left(\frac{x_1 - z}{x_0} \right) \lambda(Y_1^*(Y(v, x_3)u, x_1)w_{(1)} \otimes w_{(2)}) \\
& \left. - \operatorname{Res}_{x_3} x^{-1} \delta \left(\frac{x_0 + x_3}{x} \right) \cdot \right. \\
& \left. \cdot x_0^{-1} \delta \left(\frac{z - x_1}{-x_0} \right) \lambda(w_{(1)} \otimes Y_2(Y(v, x_3)u, x_1)w_{(2)}) \right). \quad (11.5)
\end{aligned}$$

Using (5.2) and (5.4), we find that the right-hand side of (11.5) becomes

$$\begin{aligned}
& \left(Y'_{Q(z)}(v, x) \tau_{Q(z)} \left(z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y_t(u, x_0) \right) \lambda \right) (w_{(1)} \otimes w_{(2)}) \\
& - \left(\operatorname{Res}_{x_3} x_0^{-1} \delta \left(\frac{x - x_3}{x_0} \right) \cdot \right. \\
& \left. \cdot \tau_{Q(z)} \left(z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) Y_t(Y(v, x_3)u, x_0) \right) \lambda \right) (w_{(1)} \otimes w_{(2)}). \quad (11.6)
\end{aligned}$$

By the compatibility condition for λ and the commutator formula for $Y'_{Q(z)}$, the right-hand side of (11.6) is equal to

$$\begin{aligned}
& z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) (Y'_{Q(z)}(v, x) Y'_{Q(z)}(u, x_0) \lambda) (w_{(1)} \otimes w_{(2)}) \\
& - z^{-1} \delta \left(\frac{x_1 - x_0}{z} \right) \left(\operatorname{Res}_{x_3} x_0^{-1} \delta \left(\frac{x - x_3}{x_0} \right) \cdot \right.
\end{aligned}$$

$$\begin{aligned}
& \cdot Y'_{Q(z)}(Y(v, x_3)u, x_0)\lambda)(w_{(1)} \otimes w_{(2)}) \\
&= z^{-1} \delta\left(\frac{x_1 - x_0}{z}\right) \left(\left(Y'_{Q(z)}(v, x) Y'_{Q(z)}(u, x_0) \right. \right. \\
&\quad \left. \left. - \operatorname{Res}_{x_3} x_0^{-1} \delta\left(\frac{x - x_3}{x_0}\right) Y'_{Q(z)}(Y(v, x_3)u, x_0) \right) \lambda \right) (w_{(1)} \otimes w_{(2)}) \\
&= z^{-1} \delta\left(\frac{x_1 - x_0}{z}\right) (Y'_{Q(z)}(u, x_0) Y'_{Q(z)}(v, x) \lambda) (w_{(1)} \otimes w_{(2)}). \tag{11.7}
\end{aligned}$$

This proves (11.1) and hence the first part of Proposition 6.2.

Finally, suppose that $\lambda \in (W_1 \otimes W_2)^*$ satisfies the local grading-restriction condition. We want to prove that for any $v \in V$, the components of

$$Y'_{Q(z)}(v, x)\lambda = \sum_{n \in \mathbb{Z}} \tau_{Q(z)}(v \otimes t^n) \lambda x^{-n-1}$$

also satisfy the local grading-restriction condition. From (9.6), for $n \in \mathbb{Z}$ the coefficient $\tau_{Q(z)}(v \otimes t^n)$ of x^{-n-1} in $Y'_{Q(z)}(v, x)$ has weight $\operatorname{wt} v - n - 1$, that is, $\tau_{Q(z)}(v \otimes t^n)$ maps a weight vector of weight h to a weight vector of weight $\operatorname{wt} v - n - 1 + h$. Thus the components of $Y'_{Q(z)}(v, x)\lambda$ are sums of weight vectors since λ is a sum of weight vectors. Now note that for any $v \in V$ and $n \in \mathbb{Z}$, the (graded) subspace $W_{\tau_{Q(z)}(v \otimes t^n)\lambda}$ (recall the notation in part (b) of the local grading-restriction condition) is contained in the subspace W_λ and since W_λ satisfies (6.2) and (6.3), so must $W_{\tau_{Q(z)}(v \otimes t^n)\lambda}$. Thus $\tau_{Q(z)}(v \otimes t^n)\lambda$ satisfies the local grading-restriction condition for all $v \in V$ and $n \in \mathbb{Z}$, and the second and remaining part of Proposition 6.2 is proved.

References

- [FHL] I. B. Frenkel, Y.-Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, preprint, 1989; *Memoirs Amer. Math. Soc.*, Vol. 104, Number 494, Amer. Math. Soc., Providence, 1993.
- [HL] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, I, *Selecta Mathematica*, to appear.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, PA 19104

Current address: Department of Mathematics, Rutgers University, New Brunswick, NJ 08903

E-mail address: yzhuang@math.rutgers.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08903

E-mail address: lepowsky@math.rutgers.edu